

PERLANE POST PROCEDURE INSTRUCTIONS

- Immediately after the treatment with Perlane there may be slight redness, swelling, tenderness and an itching sensation in the treated area. This is a normal result of the Perlane injection. If the inconvenience continues or if other reactions occur, please contact the office.
- Sleeping with your head elevated may help with initial swelling.
- The initial swelling after the Perlane treatment may last longer than other filler products. Some patients experience swelling for about one week. This means that the result directly after the Perlane treatment should not be seen as the final result.
- Avoid touching the Perlane treated area 6 hours after the treatment. After that the Perlane treated area can be gently washed with soap and water and light make up can be applied.
- Until the initial swelling and redness have resolved, do not expose the Perlane treated area to intense heat or extreme cold.
- Notify Dr. Chilukuri if you have persistent swelling, redness or itching lasting more than 48 hours, or if any crust, scab, or rash develops.
- If you have any significant bruising from your procedure, please call Dr. Chilukuri, he can utilize our laser to accelerate the resolution of the bruising, at no additional cost.

Perlane contains the body's own natural material, hyaluronic acid.

We want you to have an outstanding result. If you have any questions or unexpected concerns, please call the office for assistance.

During Office Hours
713.344.0450

After Hours (Cell)
713.443.8731